

FAST ALGORITHMIC NIELSEN-THURSTON CLASSIFICATION OF FOUR-STRAND BRAIDS

MATTHIEU CALVEZ AND BERT WIEST

ABSTRACT. We give an algorithm which decides the Nielsen-Thurston type of a given four-strand braid. The complexity of our algorithm is quadratic with respect to word length. The proof of its validity is based on a result which states that for a reducible 4-braid which is as short as possible within its conjugacy class (short in the sense of Garside), reducing curves surrounding three punctures must be round or almost round. As an application, we give a polynomial time solution to the conjugacy problem for non-pseudo-Anosov four-strand braids.

1. INTRODUCTION

1.1. Statement of the main result. In the 1980's Thurston gave a complete classification of the elements of the mapping class groups of surfaces: periodic, pseudo-Anosov, or reducible.

During the 1990's, algorithms which decide the Nielsen-Thurston type of a given mapping-class were constructed via the theory of train-tracks ([2], [16]). Unfortunately, the complexity of these algorithms remains unknown. Even in the particular case of the n -braid group B_n (i.e. the mapping class group of an n -times punctured disk) the general problem of deciding whether a given braid is reducible or not (which we call *reducibility problem*) has currently no known polynomial solution.

An alternative algorithm in this particular case, using Garside theory, is given in the paper [14] (which builds on [15]). However, the complexity of the algorithm in [14], while conjectured to be polynomial, strongly depends on an open question. In the present paper we give a polynomial solution to the reducibility problem in the particular case of 4-braids. More precisely we establish the following result:

Theorem 1.1. *There is an algorithm which decides the Nielsen-Thurston type of any given 4-braid x , and whose running time is $O(\mathfrak{l}^2)$, where \mathfrak{l} denotes the length of x in the classical Artin generators σ_i .*

Corollary 1.2. *There is a polynomial time solution to the conjugacy search problem for 4-strand braids in the non-pseudo-Anosov case, i.e. deciding whether two braids, at least one of which is not pA , are conjugate, and if they are, finding a conjugating element.*

The plan of the paper is as follows. In this first section we recall some facts about reducible braids and Garside theory; the second section is devoted to the proof of

Theorem 1.1 and Corollary 1.2, modulo the key technical result (Proposition 1.9) whose proof is deferred to the third section. Finally in the fourth section we give some examples and conjectures related to the reducibility problem in braid groups.

1.2. Reducible braids. Let D_n be the closed disk in \mathbb{C} with diameter $[0, n + 1]$ and with the points $\{1, \dots, n\}$ removed. It is known that the n -braid group B_n is identified with the mapping class group of D_n .

Hence there is a (right) action of the braid group on the set of isotopy classes of simple closed curves in D_n . By abuse of notation we do not distinguish between a simple closed curve and its isotopy class. We denote the curve resulting from the action of the braid x on the curve \mathcal{C} by $\mathcal{C} * x$. A simple closed curve is said to be nondegenerate if it surrounds more than one puncture and less than n .

A braid x is said to be *reducible* if it preserves setwise a family of nondegenerate simple closed curves; such a curve is then called a reduction curve for x . A reduction curve of x is said to be *essential* if it does not cross any other reduction curve. The set of all essential reduction curves of x is called the *canonical reduction system* of x and denoted by $CRS(x)$. It is known that the set $CRS(x)$ is non-empty if and only if x is reducible nonperiodic (see [6]).

A braid x is said to be *periodic* if some power of x is a power of the full twist Δ^2 . Pure periodic braids are known to be powers of Δ^2 (where Δ is the half-twist of all strands, defined in Artin's generators by the formula

$$\Delta = (\sigma_1 \dots \sigma_{n-1})(\sigma_1 \dots \sigma_{n-2}) \dots (\sigma_1 \sigma_2) \sigma_1.$$

In what follows we will take “reducible” to mean “reducible nonperiodic”. Note that a braid x is reducible if and only if every power x^t (with $t \neq 0$) of x is reducible. Note also that reducibility is a property invariant under conjugation.

The following definition, which comes from [18], uses the notion of canonical length of a braid, which will be recalled in the next subsection.

Definition 1.3. *We say that a simple closed curve in D_n is round if it is homotopic to a geometric circle. The complexity of a simple closed curve \mathcal{C} in D_n is defined to be the smallest canonical length of a positive braid which sends \mathcal{C} to a round curve (or equivalently which sends a round curve to \mathcal{C}).*

Hence the curves of complexity 0 are the round curves; we shall call *almost-round* the curves of complexity 1. In Figure 1 are represented three simple closed curves in D_4 : the first is round, the second is almost-round and is sent to a round curve by the permutation braid $\sigma_1 \sigma_3$, the third is of complexity 2 and is obtained from the first by applying σ_2^{-2} .

Let us introduce a notion which is closely related to Definition 1.3 (see [15]): the minimal standardizer of a family \mathcal{C} of disjoint simple closed curves in D_n is defined to be the smallest positive braid (for the prefix order on B_n) which sends \mathcal{C} to a family of round curves. Hence the complexity of a curve \mathcal{C} coincides with the canonical length of its minimal standardizer. In particular, we may consider the minimal standardizer of the canonical reduction system of a reducible braid.




FIGURE 1

1.3. Garside theory. For an introduction to the classical Garside structure on braid groups, the reader is referred to the papers [9] and [7] where the notion of left normal form is defined. The second one also introduces the notions of \inf , \sup and canonical length. Recall that if a braid $x = \Delta^p x_1 \cdots x_r$ is in left normal form (where the x_i 's are positive permutation braids) then $\sup(x) = p + r$, $\inf(x) = p$ and the canonical length of x , denoted by $\ell(x)$, is defined by $\ell(x) = r$.

We also recall that the super summit set $SSS(x)$ of a braid x is the (finite, nonempty) subset of the conjugacy class of x containing all those elements which have both minimal \sup and maximal \inf (or equivalently minimal canonical length) within their conjugacy class [7]. The minimal value of ℓ within the conjugacy class of a braid x is called summit canonical length of x and is denoted $\ell_s(x)$.

In order to be able to construct at least one element of $SSS(x)$ for any given braid x , let us recall a special kind of conjugation, the so-called *cyclic sliding* [11]. Let us denote by τ the conjugation by the half-twist Δ , by \wedge the gcd associated to the prefix order and, for a simple element z of B_n , by $\partial(z)$ the right complement to Δ , that is $\partial(z) = z^{-1}\Delta$.

Definition 1.4 ([11]). *Let $x = \Delta^p x_1 \cdots x_r$ be a braid in left normal form, with $r > 0$; one defines the cyclic sliding, denoted \mathfrak{s} , by the following conjugation:*

$$\mathfrak{s}(x) = \mathfrak{p}(x)^{-1} x \mathfrak{p}(x) \quad \text{where } \mathfrak{p}(x) = \tau^{-p}(x_1) \wedge \partial(x_r).$$

The simple braid $\mathfrak{p}(x)$ is called the preferred prefix of x . (If $\ell(x) = 0$, one also can define cyclic sliding: in this case x is some power of Δ and \mathfrak{s} is the trivial conjugation).

In [11] it was shown that, starting from x and iteratively applying the cyclic sliding operation $(\ell(x) - 1) \cdot \binom{n(n-1)}{2} - 1$ times yields an element of $SSS(x)$. (This is analogue to previous results of [7] and [5] concerning the so-called cycling and decycling operations.) This leads to the following result:

Theorem 1.5 ([11]). *There is an algorithm which takes as its input a braid x with n strands, runs for time $O(\ell(x)^2 n^2)$, and outputs an element x' of $SSS(x)$ which is of the form $x' = \mathfrak{s}^k(x)$ for some $k \in \mathbb{N}$.*

The cyclic sliding operation behaves well with respect to the reducibility problem:

Proposition 1.6 ([12]). *Let $x = \Delta^p x_1 \cdots x_r$ be a braid in left normal form and let \mathcal{C} be a round simple closed curve such that $\mathcal{C} * x = \mathcal{C}$. Then the curve $\mathcal{C} * \mathfrak{p}(x)$ is also round. In particular, the braid $\mathfrak{s}(x)$ also preserves a round curve.*

Now every reducible braid has some conjugate with round essential reduction curves. Thus Proposition 1.6 together with Theorem 1.5 yields:

Corollary 1.7 ([1]). *For any reducible braid x there exists an element of $SSS(x)$ with a round essential reduction curve.*

We shall also need the following result which essentially comes from [10].

Proposition 1.8. *Let $x = \Delta^p x_1 \cdots x_r$ be the left normal form of a reducible braid which preserves a simple closed curve \mathcal{C} of complexity s . Then the complexity of the curve $\mathcal{C} * \Delta^p x_1 \cdots x_i$ is bounded above by s for every i with $1 \leq i \leq r$.*

Proof. Notice that the case $s = 0$ in the proposition is proved in Theorem 5.7 in [1]. Let P be the minimal standardizer of $CRS(x)$. By Theorem 4.9 in [15], we have: $\inf(P^{-1}xP) \geq \inf(x)$ and $\sup(P^{-1}xP) \leq \sup(x)$. Under some minor changes, Proposition 2.1 and Corollary 2.2 in [10] assert the existence of positive braids $P_0 = \tau^p(P), P_1, \dots, P_r$ such that

$$P^{-1}xP \text{ has left normal } \Delta^p(P_0^{-1}x_1P_1) \dots (P_{r-1}^{-1}x_rP_r)$$

(with possibly some indices $0 \leq i' < i'' \leq r - 1$ such that $u_i^{-1}x_{i+1}u_{i+1} = \Delta$ for $i \leq i'$ and $u_i^{-1}x_{i+1}u_{i+1} = 1$ for $i \geq i''$) and

$$\inf(P_i) \geq \inf(P), \sup(P_i) \leq \sup(P) \text{ for } i = 1 \dots r.$$

The case $s = 0$ of the current Proposition yields the roundness of the curves

$$(\mathcal{C} * P) * (\Delta^p(P_0^{-1}x_1P_1) \dots (P_{i-1}^{-1}x_iP_i))$$

for all $i = 1, \dots, r$. The latter can now be rewritten as $(\mathcal{C} * (\Delta^p x_1 \dots x_i)) * P_i$. The complexity of the curve $\mathcal{C} * \Delta^p x_1 \dots x_i$ is thus at most $\ell(P_i) \leq \ell(P) = s$. This shows Proposition 1.8. \square

The special case $s = 0$ of Proposition 1.8 together with Corollary 1.7, means that for *some* (but not necessarily every) element of the SSS of a reducible braid, reducibility is easy to detect.

However this is not sufficient in order to obtain a polynomial-time algorithm for detecting reducibility since the size of the SSS may grow exponentially with both word length and braid index (see [4]). This difficulty does not yet appear in the case $n = 3$, as will be shown in Proposition 1.13. Our strategy for proving Theorem 1.1 is to show that also in the case of four strand braids, the reducibility is easy to see for *every* element in the SSS . The following is our main technical result:

Proposition 1.9. *Let x be a reducible 4-braid such that $x \in SSS(x)$. Suppose that x has an essential reduction curve surrounding 3 punctures. Then this curve is round or almost round.*

Notice that an essential reduction curve of a 4-braid x surrounding three punctures is fixed by x (not sent to another reduction curve). Now, let us recall from [11] a last Garside-theoretical notion:

Definition 1.10 ([11]). *Let $x = \Delta^p x_1 \cdots x_r$ be a braid in left normal form. We say that x is rigid if its preferred prefix is trivial, i.e. if the equality $\mathfrak{p}(x) = 1$ holds. That means that the pair $x_r \tau^{-p}(x_1)$ is left-weighted.*

We end this section with some results specifically concerning 3-braids. We start by recalling that in this case there are only four simple braids other than 1 and Δ (i.e. nontrivial strict prefixes of Δ), namely σ_1 , σ_2 , $\sigma_1\sigma_2$ and $\sigma_2\sigma_1$.

Proposition 1.11. *Let x be a 3-braid with $\text{inf}(x) = p$ and $\ell(x) = r$. Let $x_1, \dots, x_r \in B_3^+$ be non-trivial simple elements different from Δ . If $x = \Delta^p x_1 \cdots x_r$, then this is the left normal form of x .*

Proof. If it was not, making the pair $x_i x_{i+1}$ left-weighted (for some $1 \leq i \leq r-1$) either would create one factor Δ (contradicting $\text{inf}(x) = p$) or would decrease the number of factors (contradicting $\text{sup}(x) = r + p$), or both. \square

Proposition 1.12. *Let x be a 3-braid with $\ell_s(x) > 1$. Then $SSS(x)$ is the set of rigid conjugates of x .*

Proof. Let y be an element of $SSS(x)$, written $y = \Delta^p y_1 \dots y_r$ in left normal form, and suppose that y is non rigid. Consider the conjugate

$$z = (\tau^{-p}(y_1))^{-1} y \tau^{-p}(y_1) = \Delta^p y_2 \dots y_r \tau^{-p}(y_1)$$

of y . As in the proof of Proposition 1.11, the non left-weightedness of the pair $y_r \tau^{-p}(y_1)$ implies that z has smaller canonical length than y . This is a contradiction because $y \in SSS(x)$.

Conversely, it is easy to see, according to Theorem 1.5, that a rigid braid y conjugate to x belongs to $SSS(x)$ since in that case, $\mathfrak{s}(y) = y$. \square

Proposition 1.13. *The reducibility problem as well as the conjugacy search problem can be solved in polynomial time for three-strands braids.*

Proof. We claim that in this particular case, the size of the Super Summit Set of a braid x is bounded above by $2 \cdot \ell_s(x)$. Thus computing the whole Super Summit Set of any 3-braid is doable in polynomial time according to Proposition 6.2 in [8] and consequently, the reducibility problem and the conjugacy search problem can be solved quickly.

Let us prove the claim we made. First, SSS of elements of canonical length 0 (that is powers of Δ) contain only one element. Then, any braid of canonical length 1 is an element of its Super Summit Set since it is not conjugate to a power of Δ and thus the canonical length is already minimal within the conjugacy class. Thus as both letters σ_1 and σ_2 are conjugate to each other (by Δ) and because inf has constant value in the Super Summit Set one has

$$\begin{aligned} SSS(\Delta^k \sigma_1) &= \{\Delta^k \sigma_1, \Delta^k \sigma_2\}, \\ SSS(\Delta^k \sigma_1 \sigma_2) &= \{\Delta^k \sigma_1 \sigma_2, \Delta^k \sigma_2 \sigma_1\}, \end{aligned}$$

for any $k \in \mathbb{Z}$.

Suppose then that we are given a braid x of summit canonical length $\ell > 1$ together with an element $y = \Delta^p y_1 \dots y_\ell$ (in left normal form) of $SSS(x)$. Recall the cycling operation, which consists, for a general braid $u = \Delta^p u_1 \dots u_r$ in left normal form, in the conjugation $\mathbf{c}(u) = (\tau^{-p}(u_1))^{-1} u \tau^{-p}(u_1)$ [7]. Applied to (Proposition 1.12) our rigid braid y , we see that $\mathbf{c}(y) = \Delta^p y_2 \dots y_\ell \tau^{-p}(y_1)$ is in left normal form as written and rigid. Notice that $\tau(y)$ as well belongs to $SSS(x)$ and consider the

first ℓ cyclings of both y and $\tau(y)$. This produces at most 2ℓ elements of $SSS(x)$ (actually we have $\mathbf{c}^\ell(y) = \tau^{-p}(y)$, which is y if p is even and $\tau(y)$ if p is odd (recall that Δ^2 is central) so that we have just computed either two closed orbits under cycling conjugate by Δ to each other or one closed orbit under cycling, self conjugated by Δ). We thus have a subset \mathcal{O} of $SSS(x)$ closed by cycling and conjugation by Δ . By Proposition 4.14 in [8], any other element $v \in SSS(x)$ must be the result of the conjugation of some $z = \Delta^p z_1 \dots z_\ell \in \mathcal{O}$ by a minimal conjugator s . This minimal conjugator is a simple, and is shown in Corollary 2.7 in [3] (one can replace USS by SSS in this result without any change more) to be either a prefix of $\tau^{-p}(z_1)$ or of $\partial(z_r)$ (not of both since z is rigid); since v is rigid, one sees that s must actually be the whole factor, either $\tau^{-p}(z_1)$ (in which case $v = \mathbf{c}(z)$) or $\partial(z_\ell)$ (otherwise $s^{-1}vs$ would not be rigid). Noticing that

$$\partial(z_\ell)^{-1}z\partial(z_\ell) = \mathbf{c}^{\ell-1}(\tau^{p+1}(z))$$

shows that we already had all the elements of $SSS(x)$ in \mathcal{O} . They are at most 2ℓ , as claimed. \square

2. PROPOSITION 1.9 IMPLIES THEOREM 1.1

In this section we prove Theorem 1.1 with the aid of Proposition 1.9. The first observation is that there are only 2 round curves and 4 almost-round curves surrounding 3 punctures in D_4 (see Figure 2).




FIGURE 2. Above, the round curves surrounding 3 punctures in D_4 ; below, the set of curves obtained from the above ones by applying a simple braid; i.e. the set of almost-round curves surrounding 3 punctures.

Lemma 2.1. *There is an algorithm which decides in time $O(\ell^2)$ whether a 4-braid x given as a word of length ℓ in Artin's generators, is reducible with an essential reduction curve surrounding 3 punctures.*

Proof. We recall that $\ell(x) \leq l(x) \leq 6\ell(x)$. Now according to Theorem 1.5, an element y of $SSS(x)$ can be computed in time $O(\ell^2)$. According to Proposition 1.9, x admits an essential reduction curve surrounding 3 punctures if and only if y preserves one of the six curves in Figure 2. The check whether this is the case can be performed in time $O(\ell)$: according to Proposition 1.8, it is sufficient to test whether the images of these curves under each successive Garside factor of y are still round or almost round. \square

In order to overcome the fact that we don't have any analogue of Proposition 1.9 for curves surrounding 2 punctures, we will consider 4-braids as mapping classes of a 5-punctured sphere by collapsing the boundary of D_4 . The five punctures lie on the equator and we shall number them from 1 to 5, the fifth being the new puncture, on the far side of the sphere.

In the following construction x will be supposed to be pure; this assumption is fulfilled up to taking a power 2, 3 or 4. Then for $j = 1, \dots, 4$, blowing up the j th puncture to become the boundary of a new 4-punctured disk D_4 yields from x a new 4-braid which we will denote by \tilde{x}_j .

Lemma 2.2. *For $j = 1, \dots, 4$, the length $l(\tilde{x}_j)$ of \tilde{x}_j in Artin's generators is bounded above by $3l(x)$.*

Proof. We show that each letter of x gives rise to no more than 3 letters. Let us describe in detail how the first letter of x is transformed. In order to identify the j th puncture with the new boundary we make a rotation of the equator bringing j th puncture behind the sphere at the place of the fifth, and then we renumber the punctures following the rule:

$$\begin{aligned} i &\rightsquigarrow i - j + 5 && \text{if } i \leq j, \\ i &\rightsquigarrow i - j && \text{if } i > j. \end{aligned}$$

If the first letter of x is σ_i , for $1 \leq i \leq j - 2$ or for $j + 1 \leq i \leq n - 1$, then its image in \tilde{x}_j is easy to compute: it is σ_{i-j+5} or σ_{i-j} , respectively. If $j \geq 2$ and the first letter of x is σ_{j-1} , then this corresponds to a move of the puncture numbered $j - 1$ which goes to the right above the puncture numbered j . After our rotation the corresponding move involves the fourth puncture which goes above the other punctures to the first position. This corresponds to the braid $\sigma_3^{-1}\sigma_2^{-1}\sigma_1^{-1}$. In a similar way we can compute the images of all Artin's generators:

$$\begin{aligned} \sigma_i^{\pm 1} &\rightsquigarrow \sigma_{i-j+5}^{\pm 1} && \text{if } i < j - 1, \\ \sigma_i^{\pm 1} &\rightsquigarrow \sigma_{i-j}^{\pm 1} && \text{if } i > j, \\ \sigma_{j-1}^{\pm 1} &\rightsquigarrow \sigma_3^{\mp 1}\sigma_2^{\mp 1}\sigma_1^{\mp 1} && (\text{for } j \geq 2), \\ \sigma_j^{\pm 1} &\rightsquigarrow \sigma_1^{\mp 1}\sigma_2^{\mp 1}\sigma_3^{\mp 1} && (\text{for } j \leq 3). \end{aligned}$$

Notice that the first letter of x induces a permutation of the punctures which possibly sends the puncture numbered j to another position. Thus computing the image of the second letter of x with the aid of the above formulae requires a renumbering of the punctures, according to the permutation involved. The images of the following letters of x are computed in the same way. \square

Lemma 2.3. *Let x be a reducible braid without any essential reduction curve surrounding 3 punctures. Then there exists j between 1 and 4 such that \tilde{x}_j is reducible with an essential reduction curve surrounding 3 punctures.*

Proof. Note that the reducibility of x is equivalent to the reducibility of each \tilde{x}_j for $j = 1, \dots, 4$. Now under the assumption of the lemma, x admits an essential reduction curve surrounding 2 punctures. After collapsing the boundary of D_4 this curve divides the sphere into two connected components, one with 2 punctures,

the other with 3 punctures. Blowing up one of the first two to the new boundary achieves the proof of the lemma. \square

To conclude the proof of Theorem 1.1, we can give the algorithm which solves the reducibility problem in B_4 .

- 0 INPUT: A braid word x in the letters $\sigma_i^{\pm 1}$.
- 1 Compute a pure power x^t of x and do $x := x^t$.
- 2 Compute the left normal form of x .
- 3 Test whether x is periodic (that is test if x is a power of Δ^2). If yes, then RETURN “ x periodic” and STOP. Else go to 4.
- 4 Apply iterated cycling to x until the inf has not increased during the last five iterations. Do the same with decycling until the sup has not decreased during the five last iterations.
- 5 For the element of $SSS(x)$ obtained at the previous step, test whether it has a round or almost round essential reduction curve surrounding 3 punctures. If yes, RETURN “ x reducible” and STOP. If not go to the following step.
- 6 For $j = 1, \dots, 4$, compute \tilde{x}_j , apply to it steps 2 and 4. Test whether the element of $SSS(\tilde{x}_j)$ thus obtained has a round or almost round essential reduction curve surrounding 3 punctures; if the answer is positive for some j , RETURN “ x reducible” and STOP. If the answer is negative for all j , RETURN “ x pseudo-Anosov” and STOP.

Proof of Corollary 1.2. Given two braid words $x^{(1)}$ and $x^{(2)}$ of length l in the generators $\sigma_i^{\pm 1}$ ($i = 1, 2, 3$), we want to test whether they are conjugate, and we want to do so in time $O(P(l))$, where P is a polynomial. In order to achieve this, we first check whether they are periodic. If one of them is and the other one isn't, then they are not conjugate; if they both are, we are able to solve the conjugacy search problem for them in polynomial time, see [4].

Next we want to check whether both braids admit a reducing curve surrounding three punctures. In order to do so, we calculate (using iterated cycling) braids P_1 and P_2 such that $P_i^{-1}x^{(i)}P_i \in SSS(x^{(i)})$. For each of these conjugated braids we can decide whether they admit a round or almost round reducing curve surrounding 3 punctures (see Lemma 2.1). If one of them has and the other one has not then they are not conjugate. If they both have then, possibly after adding one factor to P_i , we can even assume that $P_i^{-1}x^{(i)}P_i$ (for $i = 1, 2$) both have *round* reducing curves. These two braids are now conjugate if and only if two conditions are satisfied. Firstly the winding numbers of the outer strand with the three inner strands must coincide, and secondly the inner 3-braids must be conjugate. Both of these conditions can be checked very quickly (see Proposition 1.13).

The most difficult case occurs when neither of the braids admit a reducing curve surrounding three punctures, and we have to check for reducing curves surrounding two punctures. If a braid $x^{(i)}$ has a canonical reducing curve surrounding two punctures, then a conjugate of $x^{(i)}$ in which the reducing curve is round can be found as follows. We consider the twelfth power $y^{(i)} = x^{(i)12}$ which is guaranteed to be pure and which has the same canonical reduction curves as $x^{(i)}$. One of the braids $\tilde{y}_j^{(i)}$ ($j = 1, \dots, 4$) has a canonical reduction curve surrounding three punctures.

As in the previous paragraph, we can find a braid \tilde{P}_i so that $\tilde{P}_i^{-1}\tilde{y}_j^{(i)}\tilde{P}_i$ has a *round* canonical reduction curve surrounding three punctures. The corresponding homeomorphism of the five times punctured sphere also has a reduction curve intersecting the equator only twice. Thus as in the proof of Lemma 2.2 we can explicitly write a 4-strand braid P_i such that the pure braid $P_i^{-1}y^{(i)}P_i$ has a round reducing curve surrounding two punctures. Thus the twelfth root $P_i^{-1}x^{(i)}P_i$ has the same round reducing curves.

To summarize, if the search for round reducing curves surrounding two punctures is unsuccessful in both braids $x^{(i)}$ ($i = 1, 2$), then both braids are pseudo-Anosov, and we cannot answer the question whether they are conjugate. If for one of them the search is successful and for the other it is not, then the two braids are not conjugate. If for both braids we find conjugates with round reducing curves surrounding two punctures, then we consider for each of them a 3-braid $z^{(i)}$ obtained by merging the two inner strands of a reducing curve into a single strand. We then solve the reducibility problem for each of the $z^{(i)}$'s. This can be done quickly, according to Proposition 1.13. Our two braids $x^{(i)}$ are conjugate only if the braids $z^{(i)}$ have the same Nielsen-Thurston type.

The braids $z^{(i)}$ cannot be periodic because otherwise the braids $x^{(i)}$ would have a reducing curve surrounding 3 punctures. If the $z^{(i)}$'s are pA, then the braids $x^{(i)}$ are conjugate if and only if the following two conditions are satisfied. Firstly, the winding numbers of the two inner strands must coincide, and secondly, the two three-strand braids $z^{(i)}$ must be conjugate. Again, both of these conditions can be checked in polynomial time. Finally if both braids $z^{(i)}$ are reducible, one can find 4-braids $u^{(i)}$ conjugate to $x^{(i)}$ so that $u^{(i)}$ has 2 round reducing curves, each of them surrounding two punctures. Moreover, the conjugating element can be explicitly written as a product of simple 4-braids corresponding to the conjugating 3-braids occurring during iterated cycling applied to $z^{(i)}$, but with the strand corresponding to the round reducing curve of $P_i^{-1}x^{(i)}P_i$ duplicated. Now, the braids $x^{(i)}$ are conjugate if and only if the winding number of the two fat strands in $u^{(1)}$ is the same as in $u^{(2)}$, and if the two winding numbers of the pairs of strands inside the two tubes of $u^{(1)}$ coincide with those in $u^{(2)}$. This can once again be checked quickly. \square

3. PROOF OF PROPOSITION 1.9

3.1. Outline of the proof, notation. We shall give a proof by contradiction. So let us suppose that there exists a reducible 4-braid in its own SSS which admits an essential reduction curve of complexity 2 or greater than 2 surrounding 3 punctures. By multiplying the braid by a sufficiently high power of Δ^2 one may suppose that it is positive. Moreover we may suppose this essential reduction curve to be of complexity exactly 2. If it was greater, then the ‘‘convexity’’ of the SSS (see Corollary 4.2. of [7]) and the existence in the SSS of a braid with round essential reduction curve would yield another braid in the SSS with a complexity 2 essential reduction curve.

Hence we start with a positive reducible 4-braid x in its own SSS with an essential reduction curve of complexity 2 surrounding 3 punctures. Let us denote this curve by \mathcal{C} . We recall that \mathcal{C} is fixed by x , because it is essential. We call *outer* the

strand whose puncture base is not surrounded by the curve \mathcal{C} , the other three are *inner*. For the rest of this section we denote by \hat{x} the 3-braid $z^{(i)}$ obtained from x by removing the outer strand.

The plan of the proof is as follows: first we list all curves of complexity 2 surrounding 3 punctures in D_4 . Next we prove that x and \hat{x} have the same canonical length, and that \hat{x} is a rigid braid. Finally, using a careful case-by-case analysis, we show that Proposition 1.8 and the rigidity of \hat{x} together imply that none of the curves in our list can be an essential reduction curve for x .

3.2. Curves of complexity 2 surrounding 3 punctures in D_4 . Let us classify the simple closed curves of complexity 2 surrounding 3 punctures in D_4 . In order to describe them we may also consider isotopy classes of smooth arcs which start at one of the punctures and end on the boundary of the disk, which moreover cross every vertical line in the disk at most once and the horizontal axis at least once (not counting the endpoints); these arcs are to be considered up to allowing the ending point to slide along the boundary of the disk. The bijective correspondence between the two notions is as follows: given such an arc γ we consider a tubular neighborhood N of $\gamma \cup \partial D_4$; the corresponding simple closed curve is then the boundary of $D_n - N$.




FIGURE 3

Figure 3 gives all the possibilities of such arcs. To see that these are the only possibilities consider for instance arcs starting at the first puncture. Unless to have

zero intersection with the horizontal axis, one can assume them (up to sliding the endpoint) to end at the rightmost point of the horizontal diameter of the disk. Now, the respective positions of the three remaining punctures, below or above the arc, give all the possibilities (the 8 obvious ones minus those 2 where all the punctures are simultaneously below or above).

We shall first make some remarks concerning this figure which will be useful for the end of this section:

- Remark 3.1.**
- 1) Curves of type 1 and 2 are symmetric to each other with respect to the horizontal axis. Hence acting by σ_i on one of these curves and by σ_i^{-1} on the other yields two curves with the same symmetry. The curves of type 3 and 4 have the same properties between them.
 - 2) One can obtain the curves 3 from the curves 2 (and 4 from 1) by applying the half twist Δ_4 . Moreover this symmetry is preserved by the respective actions of a braid x and its conjugate $\tau(x)$.

3.3. The two braids x and \hat{x} have the same canonical length. For a 4-braid y with an essential reduction curve surrounding 3 punctures let us denote by v_y the number of crossings in y where the outer strand is involved, counted with sign. This number is invariant under conjugacy. We remark that the winding number of the outer strand with each other strand is $\frac{v_y}{6}$.

Lemma 3.2. *Let y be a positive reducible 4-braid with an essential reduction curve of complexity 2 surrounding 3 punctures. Then $v_y < 3 \sup(y)$.*

Proof. Because y is a positive braid, y is a product of $\sup(y)$ simple braids. Because each simple braid contributes at most 3 to v_y , we have $v_y \leq 3 \sup(y)$. Suppose that the equality holds, then each simple braid contributes exactly 3 to v_y , so that the outer strand crosses all the inner strands in each Garside factor of y . If the outer strand is the first or the fourth then $\sup(y)$ is even because of the purity of this outer strand, and the circle surrounding the punctures 2, 3, 4, or 1, 2, 3, respectively, is preserved by y ; this is a contradiction since both of these circles cross each of the curves in Figure 3.

If the outer strand is the second or the third then the Garside factors of y other than Δ must be $\sigma_1\sigma_2\sigma_1\sigma_3\sigma_2$ and $\sigma_2\sigma_3\sigma_2\sigma_1\sigma_2$ alternately, so that by the purity of the outer strand, $\sup(y)$ is also even and the circle surrounding the punctures numbered 1 and 2 and the circle surrounding the punctures numbered 3 and 4 are preserved by y . We conclude as above. \square

We now see that removing the outer strand of x does not affect the canonical length:

Proposition 3.3. *The following equalities hold: $\inf(\hat{x}) = \inf(x)$ and $\sup(\hat{x}) = \sup(x)$.*

Proof. First notice that $\sup(\hat{x}) \leq \sup(x)$ and $\inf(\hat{x}) \geq \inf(x)$ because \hat{x} was obtained from x by removing a strand. Let P be the minimal standardizer of \mathcal{C} . By removing the outer strand in $P^{-1}xP$ we obtain a 3-braid x' , conjugate to \hat{x} : if $P' \in B_3$ is obtained from P by removing the outer strand, then $x' = P'^{-1}\hat{x}P'$. Since $P^{-1}xP$ preserves a round curve surrounding 3 punctures, its left normal form can be written (see [15]) as

$$P^{-1}xP = \langle x_0 \rangle x_1,$$

with $x_0 \in B_2$ and $x_1 \in B_3$. (This notation means that x_1 is the braid obtained from $P^{-1}xP$ by removing the outer strand and x_0 is the 2-braid obtained when considering x_1 as a fat strand.) Note that $x_1 = x'$. Hence if P'' is the 4-braid obtained from P' by adding a trivial strand in the suitable position and if we define $X = P''P^{-1}xPP''^{-1}$, then $X = \langle x_0 \rangle \hat{x}$; that is X is a conjugate of x in which we have a tubular braid equal to \hat{x} and an outer strand making $\frac{v_x}{6}$ twists around this “fat strand” (we also have $x_0 = \sigma_1^{v_x/3}$).

Now according to [15] one has

$$\sup(X) = \max(\sup(x_0), \sup(\hat{x}))$$

and because $x \in SSS(x)$ one has $\sup(X) \geq \sup(x)$.

Next we prove that $\sup(x_0) < \sup(X)$ (and in particular $\sup(X) = \sup(\hat{x})$). For if we had $\sup(X) = \sup(x_0)$, then $v_X = 3\sup(X)$ because of the roundness of the curve preserved by X ; and by conjugacy $v_x = v_X$. Now, according to Lemma 3.2 we also have $v_x < 3\sup(x)$. Hence

$$v_x < 3\sup(x) \leq 3\sup(X) = v_X = v_x,$$

which is a contradiction. The second part of the proposition follows as

$$\sup(\hat{x}) \leq \sup(x) \leq \sup(X) = \sup(\hat{x}).$$

In a similar way we now have

$$\inf(X) = \min(\inf(x_0), \inf(\hat{x})).$$

Moreover $\inf(X) < \inf(x_0)$, for if we had an equality, $\Delta^{-\inf(X)}X$ would be a split braid (see Section 6 in [15]). Noticing that (by purity of the outer strand) $\inf(X)$ is even, it would follow that $\Delta^{-\inf(X)}x$ is also split and positive (because $x \in SSS(x)$), one has $\inf(X) \leq \inf(x)$ so that by Proposition 6.2 in [15] the outermost curves in $CRS(\Delta^{-\inf(X)}x) = CRS(x)$ are round. This is impossible since the only circles which do not cross the curves in Figure 3 are inner to them. Hence

$$\inf(X) = \inf(\hat{x}) \geq \inf(x) \geq \inf(X),$$

where the last inequality holds since $x \in SSS(x)$. We finally obtain

$$\inf(X) = \inf(\hat{x}) = \inf(x).$$

Hence the proposition is shown and we remark also that $X \in SSS(x)$. \square

Corollary 3.4. *With the above notations we have $\hat{x} \in SSS(\hat{x})$.*

Proof. Let us suppose by contradiction that there exists a 3-braid \hat{z} in the conjugacy class of \hat{x} with $\inf(\hat{z}) > \inf(\hat{x})$ or $\sup(\hat{z}) < \sup(\hat{x})$. Let us also denote by \hat{y} the conjugating element, that is $\hat{z} = \hat{y}^{-1}\hat{x}\hat{y}$. Let z be the 4-braid obtained by conjugating X by \hat{y} augmented with a trivial strand in the suitable position, such that z has the same round essential reduction curve as X and $z = \langle x_0 \rangle \hat{z}$.

By the same argument using Lemma 3.2 as in the proof of Proposition 3.3 (with z playing the role of X), we have $\sup(z) = \sup(\hat{z})$ and $\inf(z) = \inf(\hat{z})$.

Now suppose that $\sup(\hat{z}) < \sup(\hat{x})$. Then

$$\sup(z) = \sup(\hat{z}) < \sup(\hat{x}) = \sup(x);$$

which is a contradiction, because the braids z and x are conjugate and x lies in its own SSS .

Similarly suppose that $\inf(\hat{z}) > \inf(\hat{x})$. Then $\inf(z) = \inf(\hat{z}) > \inf(\hat{x}) = \inf(x)$. This is again a contradiction. \square

Corollary 3.5. *The braid \hat{x} is rigid.*

Proof. See Proposition 1.12. \square

3.4. Analyzing the left normal form of x . Our strategy for proving Proposition 1.9 is to obtain a contradiction by proving the following statement:

Lemma 3.6. *None of the curves in Figure 3 can be an essential reduction curve for x .*

We do this by analyzing precisely the factors of the left normal form of x . These are composed of an inner 3-braid and possibly a move of the outer strand. Because of the equalities $\sup(x) = \sup(\hat{x})$ and $\inf(x) = \inf(\hat{x})$ (see Proposition 3.3), and according to Proposition 1.11, the factors of the left normal form of \hat{x} are exactly the inner components of the factors of the left normal form of x . Recall that the only nontrivial non- Δ simple elements in B_3 are $\sigma_1, \sigma_2, \sigma_1\sigma_2$ and $\sigma_2\sigma_1$.

The following table gives all the possibilities of non-trivial non- Δ positive simple braids in B_4 depending on the position of the outer strand (at the beginning of the braid) and on the value of the inner braid; for every possibility on the inner braid there are 4 possibilities for the corresponding braid on 4 strands. For the rest of this section we shall abbreviate “nontrivial non- Δ simple 4-braids” by “simple 4-braids” and they will be supposed positive unless otherwise stated.

Number of the outer strand Inner factor	1	2
σ_1	σ_2 $\sigma_2\sigma_1$ $\sigma_2\sigma_1\sigma_2$ $\sigma_2\sigma_1\sigma_2\sigma_3$	$\sigma_1\sigma_2$ $\sigma_1\sigma_2\sigma_1$ $\sigma_2\sigma_1$ $\sigma_2\sigma_1\sigma_3$
$\sigma_1\sigma_2$	$\sigma_2\sigma_3$ $\sigma_2\sigma_1\sigma_3$ $\sigma_2\sigma_1\sigma_3\sigma_2$ $\sigma_2\sigma_1\sigma_3\sigma_2\sigma_3$	$\sigma_1\sigma_2\sigma_3$ $\sigma_1\sigma_2\sigma_1\sigma_3$ $\sigma_2\sigma_1\sigma_3\sigma_2$ $\sigma_2\sigma_1\sigma_3\sigma_2\sigma_3$
σ_2	σ_3 $\sigma_3\sigma_1$ $\sigma_1\sigma_2\sigma_3\sigma_2$ $\sigma_3\sigma_1\sigma_2$	σ_3 $\sigma_3\sigma_1$ $\sigma_3\sigma_2$ $\sigma_3\sigma_2\sigma_3$
$\sigma_2\sigma_1$	$\sigma_3\sigma_2\sigma_1$ $\sigma_3\sigma_1\sigma_2\sigma_1$ $\sigma_3\sigma_2$ $\sigma_1\sigma_2\sigma_3\sigma_2\sigma_1$	$\sigma_3\sigma_2\sigma_1$ $\sigma_3\sigma_1\sigma_2$ $\sigma_3\sigma_2\sigma_3\sigma_1$ $\sigma_3\sigma_1\sigma_2\sigma_1$

We saw that curves for which the outer strand is the third or the fourth are images under the half-twist Δ of curves for which the outer strand is the second or the first, respectively. If the outer strand is the third (or the fourth) then the simple 4-braids whose interior component is σ_1 , $\sigma_1\sigma_2$, σ_2 , or $\sigma_2\sigma_1$, are images under the automorphism τ of simple 4-braids with outer strand in second position (or first position respectively) and whose inner component is σ_2 , $\sigma_2\sigma_1$, σ_1 , or $\sigma_1\sigma_2$, respectively. This allows to construct the rest of the table.

3.5. Proof of Lemma 3.6. We make a constant implicit use of Proposition 1.8, asserting that if $\Delta^p x_1 \cdots x_r$ is the left normal form of x and if x preserves one of the complexity 2 curves, then for $1 \leq i \leq r$ the image of the considered curve under $\Delta^p x_1 \cdots x_i$ is again a complexity 2 (or lower than 2) curve. By the following four lemmas we are going to eliminate all types of curves depicted in Figure 3.

Lemma 3.7. *The curves of type \mathcal{T} cannot be essential reduction curves of x .*

Proof. First notice that it is sufficient to prove the claim for curves \mathcal{T}_1 and \mathcal{T}_2 (because of the symmetries between the curves).

For both curves \mathcal{T}_1 and \mathcal{T}_2 the outer strand is the first. The simple 4-braids whose outer strand is the first and whose inner component starts with the letter σ_2 or σ_1 send the curve \mathcal{T}_1 or \mathcal{T}_2 , respectively, to strictly more complex curves. By the symmetries mentioned above, the simple 4-braids whose outer strand is the fourth and whose inner component starts with the letter σ_1 or σ_2 send the curves \mathcal{T}_4 or \mathcal{T}_3 , respectively, to strictly more complex curves. Similarly, by Remark 3.1, the action of a negative simple 4-braid whose outer strand is the first and whose inner component starts with the letter σ_1^{-1} or σ_2^{-1} on the curve \mathcal{T}_1 or \mathcal{T}_2 , respectively, yields a strictly more complex curve (as does the action of a negative simple 4-braid whose outer strand is the fourth and whose inner component starts with the letter σ_2^{-1} or σ_1^{-1} on the curve \mathcal{T}_4 or \mathcal{T}_3 , respectively).

Now suppose that the curve \mathcal{T}_1 or \mathcal{T}_2 is an essential reduction curve for x .

If $\text{inf}(x)$ is even then the first factor of $\Delta^{-\text{inf}(x)} \hat{x}$ must start with the letter σ_1 (or σ_2 , respectively), by Proposition 1.8 and the previous paragraph. By the rigidity of \hat{x} the last factor of \hat{x} (which is the inner component of the last factor of x) must end with the letter σ_1 (or σ_2 , respectively). Because of the symmetries between the curves \mathcal{T}_1 and \mathcal{T}_2 mentioned above, the image of the essential reduction curve at the beginning of the last factor of x has to be of complexity greater than 2. This contradicts Proposition 1.8.

If $\text{inf}(x)$ is odd, then the essential reduction curve at the beginning of the first non- Δ factor of x is \mathcal{T}_4 or \mathcal{T}_3 , respectively. Thus the first letter of $\Delta_3^{-\text{inf}(x)} \hat{x}$ must be σ_2 , or σ_1 , respectively. And by the rigidity of \hat{x} the last letter of \hat{x} must then be σ_1 , or σ_2 , respectively. As before we obtain a contradiction by considering the symmetries with respect to the horizontal axis.

These two cases and the remark at the beginning of the proof together achieve the proof of the lemma. \square

Lemma 3.8. *The curves of type \mathcal{U} cannot be essential reduction curves for x .*

Proof. As before we prove the statement only for curves \mathcal{U}_1 and \mathcal{U}_2 . We are going to assume, for a contradiction, that one of the curves \mathcal{U}_1 or \mathcal{U}_2 is an essential reduction curve for x . Hence Proposition 1.8 forces the image of this curve under each Garside factor of x to be of complexity at most 2.

Claim 1. *Assume that the curve \mathcal{U}_1 is an essential reduction curve for x . Then the first letter of $\Delta_3^{-\text{inf}(x)}\hat{x}$ is*

$$\begin{cases} \sigma_2 & \text{if } \text{inf}(x) \text{ is even,} \\ \sigma_1 & \text{if } \text{inf}(x) \text{ is odd.} \end{cases}$$

Proof. Figure 4(a) is a guide for the proof of this claim. We are just going to use Proposition 1.8 and the left-weightedness of pairs of consecutive factors of \hat{x} .

First suppose that $\text{inf}(x)$ is even. In order to show that the first letter of $\Delta^{-\text{inf}(x)}\hat{x}$ is σ_2 , we are going to search which simple 4-braids could be the factors of the left normal form of $\Delta^{-\text{inf}(x)}x$, provided the first letter of $\Delta^{-\text{inf}(x)}\hat{x}$ is σ_1 . Among all possible simple 4-braids whose outer strand is the second and whose inner component starts with the letter σ_1 , only two do not increase the complexity of the curve \mathcal{U}_1 . These two are $\sigma_1\sigma_2$ and $\sigma_1\sigma_2\sigma_3$. We prove that neither of these two braids can be the first factor of $\Delta^{-\text{inf}(x)}x$.

The first sends \mathcal{U}_1 to \mathcal{S}_1 and the second to \mathcal{T}_1 ; their inner components are σ_1 and $\sigma_1\sigma_2$, respectively. Now \mathcal{T}_1 is sent to strictly more complex curves by simple four-braids whose inner component starts with the letter σ_2 (by Lemma 3.7), thus the only possibility left is the first.

Now the only simple 4-braids whose inner component starts with the letter σ_1 , whose outer strand is the first (as in the curve \mathcal{S}_1) and which do not increase the complexity of the curve \mathcal{S}_1 are σ_2 and $\sigma_2\sigma_3$. The first fixes this curve and its inner braid is σ_1 , the second sends \mathcal{S}_1 to \mathcal{T}_1 and its inner braid is $\sigma_1\sigma_2$; by the same argument as above this second case is impossible.

Hence we have shown that if the first letter of $\Delta^{-\text{inf}(x)}\hat{x}$ is σ_1 then the curve \mathcal{U}_1 cannot be preserved by x . This shows the first part of the claim.

Now suppose that $\text{inf}(x)$ is odd. The essential reduction curve \mathcal{U}_1 of x is transformed (after the action of an odd number of factors Δ) into the curve \mathcal{U}_4 at the beginning of the first factor of $\Delta^{-\text{inf}(x)}x$. This situation is analogous to the situation we described in the proof of the first half of the claim, up to applying τ . This proves the second part of Claim 1.

Claim 2. *Assume that the curve \mathcal{U}_2 is an essential reduction curve for x . Then the first letter of $\Delta_3^{-\text{inf}(x)}\hat{x}$ is*

$$\begin{cases} \sigma_1 & \text{if } \text{inf}(x) \text{ is even,} \\ \sigma_2 & \text{if } \text{inf}(x) \text{ is odd.} \end{cases}$$

Proof. This proof is illustrated in Figure 4(b). Again we look at which simple 4-braids can occur in the left normal form of $\Delta^{-\text{inf}(x)}x$.

First suppose that $\text{inf}(x)$ is even. Among all simple 4-braids whose interior braid starts with the letter σ_2 and whose outer strand is the second, only two do not




FIGURE 4. (a) The action on the curve \mathcal{U}_1 of simple 4-braids whose inner component starts with the letter σ_1 . (b) The action on the curve \mathcal{U}_2 of simple 4-braids whose inner component starts with the letter σ_2 . Underlined letters σ_i indicate inner braids. Bold crosses indicate curves of complexity greater than 2. We can also see in (a) the action on the curve \mathcal{S}_1 of simple 4-braids whose inner component starts with the letter σ_1 and in (b) the action on the curve \mathcal{S}_2 of simple 4-braids whose inner component starts with the letter σ_2 . This will be used in Lemma 3.9

increase the complexity of the curve \mathcal{U}_2 , namely $\sigma_2 \sigma_3 \sigma_2$ and $\sigma_2 \sigma_3 \sigma_2 \sigma_1$. The first one has as its inner braid σ_2 and it sends \mathcal{U}_2 to \mathcal{T}_3 whereas \mathcal{T}_3 is sent to strictly more complex curves by all simple 4-braids whose inner component starts with the letter σ_2 : this case cannot occur. The second one induces $\sigma_2 \sigma_1$ on the inner strands and sends \mathcal{U}_2 to \mathcal{S}_3 .

Among all suitable simple 4-braids the only ones which do not increase the complexity of the curve \mathcal{S}_3 are $\sigma_1 \sigma_3 \sigma_2 \sigma_1$ and $\sigma_1 \sigma_3 \sigma_2 \sigma_1 \sigma_3$. Their inner components are σ_1 and $\sigma_1 \sigma_2$, and they send the curve \mathcal{S}_3 to \mathcal{T}_2 and \mathcal{S}_2 , respectively. Because simple 4-braids whose inner component starts with the letter σ_1 always send \mathcal{T}_2 to strictly more complex curves, the only possibility left is the second. The following Garside factor of x is then preceded by the curve \mathcal{S}_2 and its inner braid starts with the letter σ_2 . This situation is the image under τ of the situation at the beginning of the preceding factor. Thus this third factor must be $\sigma_1 \sigma_3 \sigma_2 \sigma_1 \sigma_3$ and at its end the curve becomes \mathcal{S}_3 , a situation we already treated. Hence we see that if the

inner braid $\Delta^{-\inf(x)}\hat{x}$ starts with the letter σ_2 , then we never retrieve the curve \mathcal{U}_2 . This shows the first part of the claim.

Now, suppose that $\inf(x)$ is odd. In this situation, the odd number of factors equal to Δ transform \mathcal{U}_2 to \mathcal{U}_3 . Up to applying τ we saw that no 3-braid starting with the letter σ_1 can be the inner braid of $\Delta^{-\inf(x)}x$, hence $\Delta^{-\inf(x)}x$ has to start with the letter σ_2 . This achieves the proof of Claim 2.

Now, since \hat{x} is rigid, if \mathcal{U}_1 is preserved by x then the last letter of \hat{x} must be σ_2 . However using the symmetry with respect to the horizontal axis, the action of negative simple 4-braids whose inner component starts with the letter σ_2^{-1} on the curve \mathcal{U}_1 can be seen from the action of positive simple 4-braids whose inner component starts with the letter σ_2 on the curve \mathcal{U}_2 . So we look at the action of the reversed word $\text{rev}(\Delta^{-\inf(x)}x)$ on the curve \mathcal{U}_2 . Though this word, with separations as in the left normal form of x , is not necessarily in left normal form, the word $\text{rev}(\Delta^{-\inf(x)}\hat{x})$ is, by Proposition 1.11. We saw in the proof of Claim 2 that this action cannot yield the curves \mathcal{U}_1 or \mathcal{U}_2 . (Notice that in the proofs of the two claims we only used the left-weightedness of the pairs of consecutive factors in \hat{x} and the fact that the image of the essential reduction curve is of complexity bounded by 2 after each Garside factor of x .) Similarly if \mathcal{U}_2 is preserved by x then the last letter of \hat{x} must be σ_1 . An argument analogous to the previous one yields the desired contradiction. \square

Lemma 3.9. *The curves of type \mathcal{S} cannot be essential reduction curves for x .*

Proof. The proof is similar to the proof of the previous lemma and can be derived with the help of Figure 4. We just have to use an additional argument since in fact there exist both: factors whose inner component starts with the letter σ_1 and which preserve the complexity of the curve \mathcal{S}_1 on one hand, and factors whose inner component starts with the letter σ_2 and which preserve the complexity of the curve \mathcal{S}_2 on the other hand. However, in both cases these factors are unique and $\Delta^{-\inf(x)}x$ only consists of repetitions of the following factors: σ_2 in the first case, yielding a circle preserved by x and which crosses the curves of type \mathcal{S} (namely the circle surrounding punctures 2, 3 and 4); $\sigma_1\sigma_3\sigma_2\sigma_1\sigma_3$ in the second case, also yielding circles having non-empty intersection with the curves of type \mathcal{S} and being preserved by x (the two circles surrounding punctures 1,2 and 3 and punctures 2,3 and 4). At this point we have achieved the analogues of Claims 1 and 2 of the previous lemma; we conclude in the same way because of the rigidity of \hat{x} . \square

Lemma 3.10. *The curves of type \mathcal{R} cannot be essential reduction curves for x .*

Proof. The proof is modeled on that of Lemma 3.8. We prove the statement only for curves \mathcal{R}_1 and \mathcal{R}_2 .

Claim 3. *Assume that the curve \mathcal{R}_1 is an essential reduction curve for x . Then the first letter of $\Delta_3^{-\inf(x)}\hat{x}$ is*

$$\begin{cases} \sigma_1 & \text{if } \inf(x) \text{ is even,} \\ \sigma_2 & \text{if } \inf(x) \text{ is odd.} \end{cases}$$

Proof. See Figure 5(a).

We first suppose that $\inf(x)$ is even. The only simple 4-braids whose outer strand is numbered 1, whose inner component starts with the letter σ_2 , and which do not increase the complexity of the curve \mathcal{R}_1 when applied to it, are $\sigma_3\sigma_2\sigma_1, \sigma_3\sigma_2$ and σ_3 . The first two have $\sigma_2\sigma_1$ as their inner braid, the third has σ_2 . Since $\mathcal{R}_1 * \sigma_3\sigma_2\sigma_1 = \mathcal{U}_1$, the following Garside factor of x (whose inner component has to start with the letter σ_1) must be $\sigma_1\sigma_2$ or $\sigma_1\sigma_2\sigma_3$ as in the proof of Lemma 3.8. This implies that in this case we can never retrieve the curve \mathcal{R}_1 , nor obtain the curve \mathcal{R}_4 .

We also have $\mathcal{R}_1 * \sigma_3\sigma_2 = \mathcal{T}_2$. Because of Lemma 3.7 we can eliminate this case. Finally σ_3 (which has σ_2 as its inner braid) fixes the curve \mathcal{R}_1 . Thus the only




FIGURE 5. (a) The action on the curve \mathcal{R}_1 of simple 4-braids whose inner component starts with the letter σ_2 . (b) The action on the curve \mathcal{R}_2 of simple 4-braids whose inner component starts with the letter σ_1 . Underlined letters σ_i indicate inner factors. Bold crosses indicate curves of complexity greater than 2. Factors which immediately yield more complex curves are not represented here.

possibility left, provided the first letter of $\Delta^{-\inf(x)}\hat{x}$ is σ_2 , is that $x = \Delta^{\inf(x)}\sigma_3^p$ for some natural integer p ; in this case x preserves a round curve (the circle surrounding punctures 2, 3 and 4) whose intersection with \mathcal{R}_1 is non-empty.

Up to applying τ (for the case when $\inf(x)$ is odd) this also shows the second statement of the claim.

Claim 4. *Assume that the curve \mathcal{R}_2 is an essential reduction curve for x . Then the first letter of $\Delta_3^{-\text{inf}(x)}\hat{x}$ is*

$$\begin{cases} \sigma_2 & \text{if } \text{inf}(x) \text{ is even,} \\ \sigma_1 & \text{if } \text{inf}(x) \text{ is odd.} \end{cases}$$

Proof. See Figure 5(b). We prove the claim only for even inf , the case of odd inf is symmetric. The only simple 4-braids whose inner component starts with the letter σ_1 and which do not increase the complexity of the curve \mathcal{R}_2 are $\sigma_2\sigma_1\sigma_2$, $\sigma_2\sigma_1\sigma_2\sigma_3$ and $\sigma_2\sigma_3\sigma_1\sigma_2\sigma_3$. The first two induce σ_1 on the inner strands. They yield \mathcal{U}_3 and \mathcal{T}_4 , respectively. By Lemma 3.7 we can exclude the latter. The first case is the image under Δ of the situation we described in Lemma 3.8. In this case we can never obtain \mathcal{R}_2 or \mathcal{R}_3 .

Finally the only possibility left is $\sigma_2\sigma_3\sigma_1\sigma_2\sigma_3$ (which induces $\sigma_1\sigma_2$ as its inner braid) and one has $\mathcal{R}_2 * \sigma_2\sigma_3\sigma_1\sigma_2\sigma_3 = \mathcal{R}_3$. According to the discussion just above, the following Garside factor of x must be $\tau(\sigma_2\sigma_3\sigma_1\sigma_2\sigma_3) = \sigma_2\sigma_1\sigma_3\sigma_2\sigma_1$ (since $\mathcal{R}_3 = \mathcal{R}_2 * \Delta$). This factor has $\sigma_2\sigma_1$ as its inner component and yields the curve \mathcal{R}_2 when applied to the curve \mathcal{R}_3 . This forces the braid $\Delta^{-\text{inf}(x)}x$ to consist of a succession of the factors $\sigma_2\sigma_3\sigma_1\sigma_2\sigma_3$ and $\sigma_2\sigma_1\sigma_3\sigma_2\sigma_1$ alternately. In this case x preserves round curves (namely the two circles surrounding the punctures 1 and 2 and the punctures 3 and 4) which cross the curve \mathcal{R}_2 . This achieves the proof of the claim.

Claims 3 and 4 determine the last letter of \hat{x} because of the rigidity of \hat{x} , and we conclude exactly as we did in the previous lemmas. \square

These last four lemmas imply Lemma 3.6; since we had an exhaustive description of curves of complexity 2 in D_4 surrounding 3 punctures, Proposition 1.9 is shown.

4. EXAMPLES AND CONJECTURES

In this last section we shall see that our result is sharp in some sense. We give some examples which we obtained with the aid of [13].

The first example shows that we cannot remove the case of almost round curves in the statement of Proposition 1.9.

Example 4.1. Let us consider the braid

$$x = \sigma_1\sigma_2\sigma_3\sigma_2.\sigma_2\sigma_1\sigma_3.\sigma_3\sigma_1.\sigma_3\sigma_2\sigma_1.\sigma_1 \in B_4,$$

which is in left normal form as written. Then $x \in SSS(x)$ and x has an almost round essential reduction curve surrounding 3 punctures and no round reduction curve. See Figure 6(a).

There are similar examples for curves surrounding two punctures:

Example 4.2. Consider the 4-braid

$$y = \sigma_3\sigma_2\sigma_3.\sigma_2\sigma_3\sigma_1.\sigma_3\sigma_2.$$

Again y lies in its own SSS , has no round reduction curve, and preserves an almost round curve surrounding 2 punctures. See Figure 6(b).




FIGURE 6. The braids x and y and their essential reduction curves. The dashed lines separate Garside factors.

Now let us show that our result cannot be extended to the $n \geq 5$ case.

Example 4.3. Consider the braid

$$z = \sigma_1 \sigma_2 \sigma_1 \sigma_3 \sigma_2 \sigma_1 \cdot \sigma_1 \sigma_3 \cdot \sigma_1 \sigma_3 \sigma_2 \sigma_4 \cdot \sigma_2 \sigma_1 \sigma_4 \sigma_3 \sigma_2 \sigma_1 \cdot \sigma_1 \sigma_2 \sigma_1 \cdot \sigma_1 \sigma_2 \sigma_1 \sigma_3 \sigma_2 \cdot \sigma_3 \in B_5,$$

which is in left normal form as written. Then $z \in SSS(z)$ and z has neither round reduction curves, nor almost round reduction curves. In fact z preserves the complexity 2 curve shown in Figure 7(a). Notice that the interior braid is pseudo-Anosov.

We conjecture that our results can be improved:

Conjecture 4.4. *Proposition 1.9 can be generalized to every simple closed curve in D_4 (even those surrounding 2 punctures) using the same kind of arguments.*

However there are more curves of complexity 2 surrounding 2 punctures in D_4 than curves of complexity 2 surrounding three punctures in D_4 , thus the proof would be more difficult.

We finish by briefly looking at two conceivable alternative approaches to the reducibility problem in B_n , for every $n \in \mathbb{N}$. The first one concerns the cyclic sliding operation:

Conjecture 4.5. *There is a polynomial bound on the number of times one has to apply cyclic sliding in order to decrease the complexity of any essential reduction curve which is not round or almost round.*

Conjecture 4.5 would imply a polynomial algorithm for solving the reducibility problem in all braid groups. This is because the results in [17] imply that the complexity of any essential reduction curve of a reducible braid is linearly bounded by the length of this braid. The following example shows that the most optimistic version of Conjecture 4.5, namely that the complexity of the reduction curve decreases in every single step, is wrong.




FIGURE 7

Example 4.6. Consider the braid z in Example 4.3. After one cyclic sliding one obtains from z the braid

$$\mathfrak{s}(z) = \sigma_1\sigma_3.\sigma_1\sigma_3\sigma_2\sigma_1\sigma_4.\sigma_2\sigma_4\sigma_3\sigma_2\sigma_1.\sigma_1\sigma_2\sigma_3\sigma_2\sigma_1.\sigma_1\sigma_2\sigma_3\sigma_2.\sigma_2\sigma_1\sigma_3\sigma_2\sigma_1.\sigma_1$$

in left normal form, which still has a curve of complexity 2, shown in Figure 7(b), as an essential reduction curve.

Another approach to the reducibility problem in B_n comes from the following:

Conjecture 4.7. *Let $x = \Delta^p x_1 \cdots x_r$ be a reducible braid in left normal form such that $x \in SSS(x)$, and let \mathcal{C} be an essential reduction curve of x . Then there is some i between 1 and r such that the curve \mathcal{C} is sent to a round or almost round curve by the braid $\Delta^p x_1 \cdots x_i$.*

Note that Theorem 1.5, together with the truth of this conjecture, would yield a polynomial time algorithm for solving the reducibility problem in B_n .

The hypothesis that $x \in SSS(x)$ in the statement of Conjecture 4.7 is necessary, as the following example shows:

Example 4.8. Let $u = \sigma_1\sigma_2\sigma_1\sigma_3\sigma_2.\sigma_2\sigma_3\sigma_2\sigma_1.\sigma_1.\sigma_1\sigma_2\sigma_3.\sigma_3\sigma_2 \in B_4$. The essential reduction curve of u is of complexity 2 and it is sent to curves of complexity 2 after each Garside factor of u (see Figure 7(c)). Notice that the inner braid is also pseudo-Anosov.

Acknowledgements The first-named author's doctoral studies are supported by a grant ARED (Région Bretagne). He is also supported by a grant for international mobility from the Université Européenne de Bretagne.

REFERENCES

- [1] D. Bernardete, M. Gutierrez, Z. Nitecki, *Braids and the Nielsen-Thurston classification*, J. Knot Theory Ramif. 4 (1995), no. 4, 549-618.
- [2] M. Bestvina, M. Handel, *Train-tracks for surface homeomorphisms*, Topology 34 (1995), no. 1, 109-140.
- [3] J. Birman, V. Gebhardt, J. González-Meneses, *Conjugacy in Garside groups II: Structure of the Ultra Summit Set*, Groups, Geom. Dyn. 2 (2008), 16-31.
- [4] J. Birman, V. Gebhardt, J. González-Meneses, *Conjugacy in Garside groups III: Periodic braids*, J. Alg. 316 (2007), 746-776.
- [5] J. Birman, K. H. Ko, S. J. Lee, *The infimum, supremum, and geodesic length of a braid conjugacy class*, Adv. Math. 164 (1) (2001), 41-56.
- [6] J. Birman, A. Lubotzky, J. McCarthy, *Abelian and solvable subgroups of the mapping class groups*, Duke Math. 50 (1983), 1107-1120.
- [7] E. ElRifai, H. Morton, *Algorithms for positive braids*, Quart. J. Math. Oxford Ser. (2) 45 (1994), 479-497.
- [8] N. Franco, J. González-Meneses, *Conjugacy problem for braid groups and Garside groups*, J. Algebra 266 (1) (2003), 112-132.
- [9] F. Garside, *The braid group and other groups*, Quart. J. Math. Oxford Ser. (2) 20 (1969), 235-254.
- [10] V. Gebhardt, *A new approach to the conjugacy problem in Garside groups*, J. Algebra 292 (2005), no. 1, 282-302.
- [11] V. Gebhardt, J. González-Meneses, *The cyclic sliding operation in Garside groups*, Mathematische Zeitschrift 265 (1), (2010), 85-114.
- [12] J. González-Meneses, *On reduction curves and Garside properties of braids*, arXiv:1006.2258.
- [13] J. González-Meneses, Braiding software, available at <http://personal.us.es/meneses>.
- [14] J. González-Meneses, B. Wiest, *Reducible braids and Garside theory*, to appear.
- [15] E. K. Lee, S. J. Lee, *A Garside theoretic approach to the reducibility problem in braid groups*, J. Algebra 320 (2008), no. 2, 783-820.
- [16] J. Los, *Pseudo-Anosov maps and invariant train tracks in the disc: a finite algorithm.*, Proc. London Math. Soc. (3) 66 (1993), no. 2, 400-430.
- [17] J. Tao, *Linearly bounded conjugator property for mapping class groups*, to appear.
- [18] B. Wiest, *How to read the length of a braid from its curve diagram*, to appear in Groups, Geom. and Dyn.

MATTHIEU CALVEZ, IRMAR (UMR 6625 DU CNRS), UNIVERSITÉ DE RENNES 1, CAMPUS DE BEAULIEU, 35042 RENNES CEDEX, FRANCE

E-mail address: `matthieu.calvez@univ-rennes1.fr`

BERT WIEST, IRMAR (UMR 6625 DU CNRS), UNIVERSITÉ DE RENNES 1, CAMPUS DE BEAULIEU, 35042 RENNES CEDEX, FRANCE

E-mail address: `bertold.wiest@univ-rennes1.fr`